

You're pregnant and have COVID-19? We're here to support you

Patient Information - Maternity Services

During pregnancy, most people with COVID-19 will have a mild flu-like illness and it is unlikely to cause problems for you or your baby, especially if you have been vaccinated.


I'm pregnant and isolating at home with COVID-19, what happens now?

- » If you have self-tested at home using a RAT kit, please upload the result of this using this link: How to report your RAT results. By doing this you let the community health team know that you have tested positive so they can then get in touch with you to offer support if you do become more unwell. If you are having trouble using My Covid Record, you can report the results of your RAT over the phone: 0800 222 478 and choose option 3.
- your family doctor or a health worker should call you soon after receiving your positive result. If they do not call you, call them. If you do not have a family doctor or health worker, please call Healthline 0800 358 5453.
- » Your family doctor or health worker will ask you some questions and tell you what to look out for and how to look after yourself.
- » They may offer you some equipment (such as a pulse oximeter) to help check your COVID-19 symptoms. This equipment will be on loan at no charge.
- » Contact your lead maternity carer (LMC) to let them know you have COVID-19.
- » Unless you become very unwell, your LMC will continue to provide your pregnancy care. This may be by phone or video call.
- » Your LMC will notify the team at the hospital that you have a COVID-19 infection, to get a plan of care for you for your pregnancy if any changes to your existing plan are needed.
- » If you become very unwell with COVID-19 you will get extra care from the team at the hospital.



What are the symptoms of COVID-19?


Most people will have mild symptoms or no symptoms at all. When symptoms occur, they can include:



For most people symptoms usually only last a few days. Unless your symptoms make you feel very ill you can usually stay at home and look after yourself.

What can I do to look after myself at home?

Diarrhoea



- » Paracetamol (such as Panadol) is safe to take for fever, headaches or muscle aches be sure to follow the instructions on the packet.
- » Do not take Ibuprofen (such as Nurofen or Brufen) as this may harm your unborn baby. Many products may contain this, so check the ingredients before taking anything sold to help with colds, fevers and blocked noses.

What are the symptoms of COVID-19?

Most people will have mild symptoms or no symptoms at all. When symptoms occur, they can include:

	If you have no symptoms or your symptoms are mild continue to check THREE times a day	Call your GP or healthcare team	Call 111 for an ambulance	Call your LMC or the Birthing Suite (03) 364 4715
Breathing	You should be breathing comfortably	If your breathing is becoming more difficult	If you have severe trouble breathing, or have severe chest pain	If you have: Tummy pain, cramps or contractions Vaginal bleeding Waters leaking If you are concerned that your baby's movements are less than usual Sudden puffiness in the face Ongoing headaches Blurred vision Pain under your ribs on right side Any other concerns about your pregnancy
Oxygen level % (if you are given a pulse oximeter)	Your oxygen level should be 95-100%	If your oxygen level is 92-94%.	If your oxygen level is below 92%	
Heart rate	Your heart rate should be less than or equal to 100 beats per minute	If your heart rate is 100-119 beats per minute	If your heart rate is above 120 beats per minute	
Temperature	Your temperature should be less than or equal to 38 degrees	If your temperature is greater than 38.5 degrees, even if you have had Paracetamol	Your GP or healthcare team will advise whether you need to call 111.	
Other symptoms		Your symptoms are getting worse Your start getting better and then get worse You have symptoms of severe dehydration such as: • a very dry mouth • passing only a little urine (pee) • feeling very light-headed.	If you are very confused or not thinking clearly If you feel faint or pass out (lose consciousness).	
Some useful links	Examples of diaries you can download or you can use as a diary yourself Click on the link or copy and paste it into your browser: https://www.healthnavigator.org. nz/media/15256/14- covid-19 health and symptom diary.pdf Click on the link or copy and paste the URL into your browser: https://www.healthnavigator.org.nz/media/15256/14- covid-19 health and symptom diary.pdf or this: https://www.healthnavigator.org.nz/media/15254/141-covid-19 symptom diary.pdf		Depending on your symptoms, you may be given a pulse oximeter Click on the link or copy and paste it into you browser https://www.youtube.com/watch?v=SyhQrjYCOGs Click on the link or copy and paste it into your browser: https://www.healthnavigator.org.nz/health-a-z/t/thermometers-how-to-use-them/	

After you have recovered

- » After you have recovered from COVID-19 and have finished isolation, your LMC will continue to provide your regular check-ups before and after your baby is born.
- » If you had mild or no symptoms you will continue to receive normal care from your LMC. For those identified as having additional risk factors, you may be offered a baby wellbeing scan at around 37-38 weeks. Sometimes having COVID-19 can affect your baby's growth. It is important that you tell your LMC if your baby movements are reduced or if you have any other concerns.
- » If you were unwell with COVID-19 during your pregnancy you will be recommended some additional scans and may have a new plan of care until your baby is born. Your LMC will let you know about any changes in your pregnancy care.
- » It is important that you tell your LMC if your baby's movements are reduced.

Vaccination after your recovery

- » If you were not vaccinated before you caught COVID-19, you can have your first or second vaccination, or booster dose four weeks after you have recovered.
- » This is a useful link about vaccination during pregnancy and breastfeeding:
- » COVID-19 vaccine: Pregnancy and breastfeeding

What happens if I need to have my baby while I am still in isolation?

- your LMC will care for you when you have your baby. Please call your LMC before you go to the hospital.
- You will be able to bring one support person with you when you are in labour or having your caesarean section.
- » Make sure your support person does not have any COVID-19 symptoms. If they do, get them to use a RAT and if they are positive ask your LMC for advice.
- » Tell your support person that they will need to wear a mask.
- » If you are both well, you and your baby will stay together and can have skin-to-skin.
- » After you have had your baby, we will do our best to enable a member of your whanāu to be with you while you are in hospital.

Will my baby get COVID-19?

- » If your baby catches COVID-19, it is likely they will be mildly unwell or have no symptoms.
- » To help protect your baby from catching COVID-19:
 - Make sure that you wash your hands with soap and water for 20 seconds before you touch your baby.
 - Wear a mask when closer than 2 metres from your baby, eg. when breastfeeding.
 - Wear a mask for 7 days from when you started to have symptoms.
- » If you are worried that your baby is unwell contact your LMC or family doctor for advice.
- » Click on this link or copy and paste it into your browser to find some useful information about breastfeeding and caring for your newborn if you have COVID-19: https://www.health.govt.nz/covid-19-vaccine-pregnancy-and-breastfeeding



Can I breastfeed if I have COVID?

- » Yes, breastfeeding your baby is recommended if you have COVID-19
- » There are antibodies in your breastmilk that will help protect your baby from COVID-19.
- you can get breastfeeding and infant feeding support from your LMC and http://canbreastfeed.co.nz/ or Te Rito Ora on 0800 HelpBF (0800 435723)
- » Click here or copy and paste the link into your browser to watch a short video about vaccination, pregnancy and breastfeeding: https://www.youtube.com/watch?v=VJTJ_Im2Eok&t=100s

Feeling anxious and stressed about COVID-19?

It is normal to feel anxious, upset, worried or stressed about COVID-19:

- » You can talk to your LMC, family doctor or health worker.
- » If you feel very worried or upset and want to talk to someone you can call 0800 111 757 or text 4202 for free, anytime to speak to a trained counsellor.
- » Click on the link or copy and paste into your browser to find some helpful resources to help you look after your mental health if you have COVID: https://depression.org.nz/covid-19/covid-19/

Useful Canterbury Health phone numbers

Healthline – COVID-19	0800 358 5453	
Christchurch Women's Hospital Birthing Suite	(03) 364 4715	
Maternity Assessment unit (MAU)	(03) 378 6410	
Maternity Outpatients	(03) 364 4454	
Christchurch Women's Hospital Maternity Ward	(03) 364 4724	
Rangiora Maternity Unit	(03) 313 7299	
Lincoln Maternity Unit	(03) 325 2802	
Kaikōura Health (Te Hā o Te Ora)	(03) 319 3500	
Ashburton Maternity Unit	(03) 307 8483	
St. George's Maternity Unit	(03) 375 6020	
Plunket Line	0800 933 922	
Healthline	0800 611 116	
Emergency Care	111	

Reproduced with permission from Counties Manakau District Health Board

For more information about: hospital and specialist services, go to www.cdhb.health.nz | health and medication advice, go to www.healthinfo.org.nz

